arXiv:gr-qc/0101056v1 15 Jan 2001

The mass quantum and black hole entropy II

B. Ram

Physics Department, New Mexico State University
Las Cruces, NM 88003 USA†

and

Umrao Institute of Fundamental Research
A2/214 Janak Puri, New Delhi, 110058, India

Abstract

In [Phys. Lett. A 265 (2000) 1] a new method was given which naturally led to a quantum of mass equal to twice the Planck mass. In the present note which, for convenience, we write formally as a continuation of that paper, we show that with spin one of the mass quantum, the physical entropy of a rotating black hole is also given by the Bekenstein-Hawking formula.

PACS: 04.60.-m, 04.70.Dy

The interest in this theory [1] lies in the facts: (1) that it combines elements of general relativity with elements of quantum mechanics in an unprecedented way; (2) that it gives the correct physical result for the entropy of a black hole, namely $A/4$, which has virtually been elevated to the status of a cardinal truth.

One may observe that the quantum eigenvalue equation (8) can be obtained by simply setting $E = \ell = 0$ in Eq. (5). This is equivalent to putting the classical constants of integration E and L equal to zero in Eq. (1) and then applying the Schrödinger prescription.

Let us apply this recipe to the classical time like Kerr geodesic [2]:

$$\frac{1}{2}\dot{r}^2 - \frac{m}{r} + \frac{1}{2}(1 - E^2)(1 + a^2/r^2) + \frac{L^2}{2r^2} - \frac{m}{r^3}(L - aE)^2 = 0, \tag{28}$$

---

† Correspondence address.

where a is the angular momentum per unit mass of the source [3]. Putting $E = L = 0$ in (28), followed by the Schrödinger prescription, gives

$$\left(-\frac{1}{2r^2}\frac{d}{dr}\left(r^2\frac{d}{dr}\right)+\frac{j(j+1)}{2r^2}-\frac{m}{r}\right)R=-\frac{1}{2}R, \tag{29}$$

or

$$\left[-\frac{1}{2}\left(\frac{d^2}{dr^2}-\frac{j(j+1)}{r^2}\right)-\frac{\mu/4}{r}\right]U=-\frac{1}{2}U, \tag{30}$$

with $U = rR(r)$ and $m = \mu/4$, as the quantum equation for the *pure* mass μ. Eq. (30) is Eq. (11) with

$$\ell' = j, \quad N' = 3, \quad B = 1/2, \quad \alpha = \mu/4;$$

and hence it represents a *four dimensional harmonic oscillator* with $\omega = 2$, and

$$\mu_{n,j} = 2(n+j+1)\omega, \qquad n = 0, 1, 2, \ldots \tag{31}$$

and the values of j determined as follows:

In modern language Eq. (31) says that the nth mass (μ) states are occupied by n *pairs* of mass quanta and $2j$ is the spin angular momentum of the n pairs.

As $n = 1$ states have only one pair, $2j$ is the spin (for brevity) of the pair. Let us designate by j_q the *spin* of the mass quantum, which can have only two values along the z-axis [3], $+j_q$ (up) and $-j_q$ (down). In principle j_q can have any positive integral value. We choose $j_q = 1$, for simplicity and because this value is in accord with the quantum version of the censorship hypothesis [4], namely

$$\sqrt{j_q(j_q+1)} \le 2 = \text{ mass of the quantum.}$$

Now the pair can be formed in only two ways: (i) when the spins of the two quanta are anti-aligned; and (ii) when the spins of the two quanta are aligned *up*. In case (i) the spin of the pair is zero, giving $2j = 0$, or $j = 0$; in case (ii) the spin of the pair is two, giving $2j = 2$, or $j = 1$. (The value $j = -1$ is not allowed because in Eq. (30) the quantum number j is a positive integer or zero.)

Next consider the $n = 2$ states. They have two pairs. These can be formed in three ways. (1) Both pairs have spin zero, giving $j = 0$; (2) one pair has spin zero and the other has spin 2, giving $j = 1$; (3) each pair has spin 2, giving $2j = 4$, or $j = 2$.

Continuing the process leads to the values $j = 0, 1, 2, \ldots, n$ for the nth mass states which contain n pairs.

Clearly then Eq. (31) can be rewritten as

$$\mu_{n'_{n,j}} = 2(n'_{n,j}+1)\omega, \qquad n'_{n,j} = 0, 1, 2, \ldots \tag{32}$$

It is to be noted that Eq. (32) is Eq. (9′) when every one of the n pairs has spin zero.

It follows straightforwardly from Eq. (32) that the physical entropy of a non-extremal Kerr black hole of mass M and angular momentum a should be exactly given by the relation (27) with $A = 16\pi M_{ir}^2$, provided the physical temperature of the black hole is defined by

$$T = \frac{\kappa}{2\pi(1-a^2/M^2)^{1/2}} = \frac{\kappa'}{2\pi}. \tag{33}$$

In conclusion we may say that the two parameters that characterize the Kerr solution of classical general relativity are fused by quantum mechanics into a single concept of a quantum of mass *twice* the Planck mass and spin angular momentum *one*.

Acknowledgements

The author thanks A. Ram, N. Banerjee, R.S. Bhalerao, and A. Bussian for useful conversations, V. Singh for affirming comments, N. Ram for critical reading of the manuscript and making concrete suggestions, and the Tata Institute for a pleasant stay.

References

[1] B. Ram, Phys. Lett. A 265 (2000) 1 [gr-qc/9908036].

[2] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, Oxford University Press, 1983.

[3] R.P. Kerr, Phys. Rev. Lett. 11 (1963) 237.

[4] B. Carter, in *Black Holes*, Editors C. DeWitt and B.S. DeWitt, Gordon and Breach, 1972.